

THE FIXED SUBGROUPS OF HOMEOMORPHISMS OF SEIFERT MANIFOLDS

QIANG ZHANG

ABSTRACT. Let M be a compact connected orientable Seifert manifold with hyperbolic orbifold B_M , and $f_\pi : \pi_1(M) \rightarrow \pi_1(M)$ be an automorphism induced by an orientation-reversing homeomorphism f of M . We give a bound on the rank of the fixed subgroup of f_π , namely, $\text{rankFix}(f_\pi) < 2\text{rank}\pi_1(M)$, which is similar to the inequalities on surface groups and hyperbolic 3-manifold groups.

1. INTRODUCTION

For a group G and an endomorphism $\phi : G \rightarrow G$, the *fixed subgroup* of ϕ is

$$\text{Fix}(\phi) := \{g \in G \mid \phi(g) = g\},$$

which is a subgroup of G . Let $\text{rank}G$ denote the *rank* of G , which means the minimal number of the generators of G .

For a free group F and an automorphism ϕ , M. Bestvina and M. Handel [BH] solved the Scott conjecture:

Theorem 1.1 (Bestvina-Handel). *Let F be a free group and ϕ be an automorphism of F . Then*

$$\text{rankFix}(\phi) \leq \text{rank}F.$$

In the paper by B. Jiang, S. Wang and Q. Zhang [JWZ], it is proved that

Theorem 1.2 (Jiang-Wang-Zhang). *Let S be a compact surface and ϕ be an endomorphism of $\pi_1(S)$. Then*

$$\text{rankFix}(\phi) \leq \text{rank}\pi_1(S).$$

In a recent paper [WZ], J. Wu and Q. Zhang generalized Theorem 1.2 to a family endomorphisms of a surface group. In [LW] (see [Z2] for an enhance version), J. Lin and S. Wang showed that

Theorem 1.3 (Lin-Wang). *Let M be a compact orientable hyperbolic 3-manifold with finite volume and ϕ be an automorphism of $\pi_1(M)$. Then*

$$\text{rankFix}(\phi) < 2\text{rank}\pi_1(M).$$

In this paper, we consider the fixed subgroups of automorphisms of Seifert 3-manifold groups.

Suppose M is a compact orientable 3-manifold. We say that M is a *Seifert manifold*, if M possesses a *Seifert fibration* which is a decomposition of M into disjoint simple closed curves, called *fibers*, such that each fiber has a solid torus neighborhood consisting of a union of fibers. Identifying each fiber of M to a point, we get a set B_M , called the *orbifold* of M , which has a natural 2-orbifold structure with singular points consisting of cone points. It is useful to think of a Seifert manifold as a circle bundle over a 2-orbifold. For brevity, we say an orbifold it means a compact 2-orbifold with singular points consisting of cone points in the following. An orbifold (or

1991 *Mathematics Subject Classification.* 57M05, 57M07, 37C25.

Key words and phrases. Seifert manifolds, fixed subgroups, homeomorphisms, ranks.

surface) is called *hyperbolic* if it has negative Euler characteristics. A hyperbolic orbifold is orbifold covered by a hyperbolic surface and admits a hyperbolic structure with totally geodesic boundary. For more information about orbifolds, see [JWW, §1 and §2] and [S, §2]. A map f on a Seifert manifold M is called *fiber-preserving* if it maps fibers to fibers. If f is fiber-preserving, then it induces a map $f' : B_M \rightarrow B_M$ on the orbifold B_M .

In the following, all spaces are assumed to be connected and compact unless it is specially stated otherwise. For a set X , let $\#X$ denote the number of points in X .

The main result of this paper is

Theorem 1.4. *Suppose M is a compact connected orientable Seifert manifold (closed or with toroidal boundary) with hyperbolic orbifold B_M , and $f_\pi : \pi_1(M) \rightarrow \pi_1(M)$ is an automorphism induced by an orientation-reversing homeomorphism $f : M \rightarrow M$. Then*

$$\text{rankFix}(f_\pi) < 2\text{rank}\pi_1(M).$$

Remark 1.5. By the well known Geometrization Theorem, M is a Seifert manifold with hyperbolic orbifold B_M if and only if M admits a geometric structure based on one of the two geometries : $\mathbb{H}^2 \times \mathbb{R}$, $SL(2, \mathbb{R})$. The condition that f is orientation-reversing is necessary. If f is orientation-preserving, then the fixed subgroup $\text{Fix}(f_\pi)$ can be infinitely generated, see [Z1, Example 5.2].

Remark 1.6. By [Z1, Example 5.1], one can show that there is an orientation-reversing homeomorphism f of $M_n = S_n \times S^1$, where S_n is a closed orientable surface of genus $n \geq 2$, such that $\text{rankFix}(f_\pi) = 4n - 2$. Therefore, for any $\varepsilon > 0$, there exists a Seifert manifold M_n and an orientation-reversing homeomorphism f of M_n , such that

$$\frac{\text{rankFix}(f_\pi)}{\text{rank}\pi_1(M_n)} = \frac{4n - 2}{2n + 1} > 2 - \varepsilon.$$

Theorem 1.4 is inspired by the following proposition [Z1, Corollary 1.4].

Proposition 1.7. *Suppose $f : M \rightarrow M$ is a homeomorphism of a compact connected orientable Seifert manifold with hyperbolic orbifold B_M . Let $f_* : \pi_1(M, x) \rightarrow \pi_1(M, x)$ be the induced automorphism, where x is a fixed point contained in an essential fixed point class of f . Then*

$$\text{rankFix}(f_*) < 2\text{rank}\pi_1(M).$$

The paper is organized as follows. In Section 2, we will give some background on fixed points and fixed subgroups of a selfmap. In Section 3, we will give some useful facts on Fuchsian groups. In Section 4, we will consider the special case of Theorem 1.4 that M is an orientable circle bundle over an orientable hyperbolic surface. Finally, we will finish the proof of Theorem 1.4 in Section 5, and give some examples and questions in Section 6.

Acknowledgements. The author would like to thank Professor Boju Jiang for valuable communications. This work was carried out while the author was visiting Princeton University and he would like to thank for their hospitality. The author is partially supported by NSFC (No. 11201364) and “the Fundamental Research Funds for the Central Universities”.

2. FIXED POINTS AND FIXED SUBGROUPS

Let X be a connected compact polyhedron and $f : X \rightarrow X$ be a selfmap. In this section, we introduce some facts on fixed point classes and fixed subgroups of f .

The fixed point set

$$\text{Fix}f := \{x \in X | f(x) = x\}$$

splits into a disjoint union of *fixed point classes*: two fixed points are in the same class if and only if they can be joined by a *Nielsen path*, which is a path homotopic (rel. endpoints) to its own f -image. For each fixed point class \mathbf{F} , a homotopy invariant *index* $\text{ind}(f, \mathbf{F}) \in \mathbb{Z}$ is defined. A fixed point class is *essential* if its index is non-zero, otherwise, called *inessential* (see [J] for an introduction).

Although there are several approaches to define fixed point classes, we state the one using paths and introduce another homotopy invariant *rank* $\text{rank}(f, \mathbf{F}) \in \mathbb{Z}$ for each fixed point class \mathbf{F} (see [JWZ, §2]), which plays a key role in this paper.

Definition 2.1. By an f -route we mean a homotopy class (rel. endpoints) of path $w : I \rightarrow X$ from a point $x \in X$ to $f(x)$. For brevity we shall often say the path w (in place of the path class $[w]$) is an f -route at $x = w(0)$. An f -route w gives rise to an endomorphism

$$f_w : \pi_1(X, x) \rightarrow \pi_1(X, x), [a] \mapsto [w(f \circ a)\overline{w}]$$

where a is any loop based at x , and \overline{w} denotes the reverse of w . For brevity, we will write

$$f_\pi : \pi_1(X) \rightarrow \pi_1(X)$$

when w and the base point x are omitted.

Two f -routes $[w]$ and $[w']$ are *conjugate* if there is a path $q : I \rightarrow X$ from $x = w(0)$ to $x' = w'(0)$ such that $[w'] = [\overline{q}w(f \circ q)]$, that is w' and $\overline{q}w(f \circ q)$ homotopic rel. endpoints. We also say that the (possibly tightened) f -route $\overline{q}w(f \circ q)$ is obtained from w by an f -route move along the path q .

Note that a constant f -route w corresponds to a fixed point $x = w(0) = w(1)$ of f , and the endomorphism f_w becomes the usual

$$f_* : \pi_1(X, x) \rightarrow \pi_1(X, x), [a] \mapsto [f \circ a],$$

where a is any loop based at x . Two constant f -routes are conjugate if and only if the corresponding fixed points can be joined by a Nielsen path. This gives the following definition.

Definition 2.2. With an f -route w (more precisely, with its conjugacy class) we associate a *fixed point class* \mathbf{F}_w of f , which consists of the fixed points that correspond to constant f -routes conjugate to w . Thus fixed point classes are associated bijectively with conjugacy classes of f -routes. A fixed point class \mathbf{F}_w can be empty if there is no constant f -route conjugate to w . Empty fixed point classes are inessential and distinguished by their associated route conjugacy classes.

Definition 2.3. The *fixed subgroup* of the endomorphism f_w is the subgroup

$$\text{Fix}(f_w) := \{\gamma \in \pi_1(X, w(0)) \mid f_w(\gamma) = \gamma\}.$$

The *stabilizer* of the fixed point class \mathbf{F}_w is defined to be

$$\text{Stab}(f, \mathbf{F}_w) := \text{Fix}(f_w),$$

it is well defined up to isomorphism because conjugate f -routes have isomorphic stabilizers. Hence, we have the *rank* of \mathbf{F}_w defined as

$$\text{rank}(f, \mathbf{F}_w) := \text{rankStab}(f, \mathbf{F}_w) = \text{rankFix}(f_w).$$

The following are some facts on stabilizer (see [JWZ, §2]).

Fact (Homotopy invariance). A homotopy $H = \{h_t\}_{t \in I} : X \rightarrow X$ gives rise to a bijective correspondence $H : \mathbf{F}_{w_0} \mapsto \mathbf{F}_{w_1}$ from h_0 -fixed point classes to h_1 -fixed point classes, and

$$\text{ind}(h_0, \mathbf{F}_{w_0}) = \text{ind}(h_1, \mathbf{F}_{w_1}), \text{Stab}(h_0, \mathbf{F}_{w_0}) \cong \text{Stab}(h_1, \mathbf{F}_{w_1}),$$

which indicates that the index $\text{ind}(f, \mathbf{F}_w)$ and the rank $\text{rank}(f, \mathbf{F}_w)$ of a fixed point class are both homotopy invariants.

Fact (Morphism). A morphism from a selfmap $f : X \rightarrow X$ to a selfmap $g : Y \rightarrow Y$ means a map $h : X \rightarrow Y$ such that $h \circ f = g \circ h$. It induces a natural function $w \mapsto h \circ w$ from f -routes to g -routes and a function $\mathbf{F}_w \mapsto \mathbf{F}_{h \circ w}$ from f -fixed point classes to g -fixed point classes, such that

$$h(\mathbf{F}_w) \subseteq \mathbf{F}_{h \circ w}, \quad h_* \text{Stab}(f, \mathbf{F}_w) \leq \text{Stab}(g, \mathbf{F}_{h \circ w}).$$

The following are useful lemmas on covering spaces and fixed point classes.

Lemma 2.4. *Let $p : \widetilde{M} \rightarrow M$ be a finite covering of a compact manifold M , and $f : M \rightarrow M$ be a homeomorphism. Suppose $\tilde{f} : \widetilde{M} \rightarrow \widetilde{M}$ is a lifting of f , and the \tilde{f} -route \tilde{w} is a lifting of the f -route w . Then the f -fixed point class \mathbf{F}_w associated to w is essential if and only if the \tilde{f} -fixed point class $\mathbf{F}_{\tilde{w}}$ associated to \tilde{w} is essential, moreover,*

$$\text{ind}(\tilde{f}, \mathbf{F}_{\tilde{w}}) = n \times \text{ind}(f, \mathbf{F}_w)$$

where n is a positive integer.

Proof. Since indices of fixed point classes are homotopy invariants of the map f , via perturbation, we may assume that all fixed points of f are isolated.

Firstly, we claim that $\mathbf{F}_{\tilde{w}}$ is nonempty if and only if \mathbf{F}_w is nonempty, and $p(\mathbf{F}_{\tilde{w}}) = \mathbf{F}_w$.

If $\mathbf{F}_{\tilde{w}}$ is nonempty, then for any point $\tilde{x} \in \mathbf{F}_{\tilde{w}} \subseteq \text{Fix} \tilde{f}$, the \tilde{f} -route \tilde{w} is conjugate to \tilde{x} , namely, there is a path $\gamma : I \rightarrow \widetilde{M}$ from $\tilde{w}(0)$ to \tilde{x} such that $\overline{\gamma} \tilde{w}(\tilde{f} \circ \gamma) \simeq \tilde{x}$ rel. \tilde{x} . Hence

$$(\overline{p \circ \gamma})w(f(p \circ \gamma)) = p(\overline{\gamma} \tilde{w}(\tilde{f} \circ \gamma)) \simeq p(\tilde{x}).$$

Namely, the f -route w is conjugate to the point $p(\tilde{x}) \in \mathbf{F}_w \subseteq \text{Fix} f$. So $p(\mathbf{F}_{\tilde{w}}) \subseteq \mathbf{F}_w$.

If \mathbf{F}_w is nonempty, then for any point $x \in \mathbf{F}_w \subseteq \text{Fix} f$, the f -route w is conjugate to x , namely, there is a path c from $w(0)$ to x such that $\overline{c}w(f \circ c) \simeq x$ rel. x . Let \tilde{c} be a lifting of c from $\tilde{w}(0)$ to a point $\tilde{x} \in p^{-1}(x)$. Then $\overline{\tilde{c}}\tilde{w}(\tilde{f} \circ \tilde{c})$ is a lifting of the contractible loop $\overline{c}w(f \circ c)$. Hence $\overline{\tilde{c}}\tilde{w}(\tilde{f} \circ \tilde{c})$ is also a contractible loop and $\tilde{f}(\tilde{x}) = \tilde{x}$, namely, the \tilde{f} -route \tilde{w} is conjugate to the point $\tilde{x} \in \mathbf{F}_{\tilde{w}} \subseteq \text{Fix} \tilde{f}$. So $\mathbf{F}_w \subseteq p(\mathbf{F}_{\tilde{w}})$ and the claim holds.

Secondly, we prove $\text{ind}(\tilde{f}, \mathbf{F}_{\tilde{w}}) = n \times \text{ind}(f, \mathbf{F}_w)$.

If \mathbf{F}_w is empty, then $\mathbf{F}_{\tilde{w}}$ is also empty according to the claim above, and the equation holds clearly.

Now we consider the case that \mathbf{F}_w is nonempty. Pick a point $x \in \mathbf{F}_w$. We can assume

$$p^{-1}(x) \cap \mathbf{F}_{\tilde{w}} = \{\tilde{x}_1, \dots, \tilde{x}_n\}$$

where $n > 0$ since the covering $p : \widetilde{M} \rightarrow M$ is finite and $p(\mathbf{F}_{\tilde{w}}) = \mathbf{F}_w$. If $y \neq x$ and $y \in \mathbf{F}_w$, then there is a Nielsen path c from x to y such that $c \simeq f \circ c$ rel. $\{x, y\}$ by the definition of fixed point class. Hence there are n liftings \tilde{c}_i with $\tilde{c}_i(0) = \tilde{x}_i$ and $\tilde{f} \circ \tilde{c}_i \simeq \tilde{c}_i$ rel. endpoints, $i = 1, \dots, n$. Therefor

$$\{\tilde{c}_1(1), \dots, \tilde{c}_n(1)\} = p^{-1}(y) \cap \mathbf{F}_{\tilde{w}}.$$

Then

$$\#\mathbf{F}_{\tilde{w}} = n\#\mathbf{F}_w.$$

So the equality holds by the fact $\text{ind}(\tilde{f}, \tilde{x}) = \text{ind}(f, x)$. \square

A covering $p : \widetilde{M} \rightarrow M$ between compact manifolds is called *characteristic* if the subgroup $p_*\pi_1(\widetilde{M})$ is a characteristic subgroup of $\pi_1(M)$, i.e., it is invariant under any automorphism of $\pi_1(M)$. Recall that if G is a finite index subgroup of a finitely generated group H , then there is a finite index characteristic subgroup G' of H , such that $G' \subseteq G$. It follows that given any finite covering $p : \widetilde{M} \rightarrow M$ of a compact manifold M , there is a finite covering $q : \widehat{M} \rightarrow \widetilde{M}$ so that $p \circ q : \widehat{M} \rightarrow M$ is a characteristic covering. In this case for any homeomorphism $f : M \rightarrow M$ with

$y = f(x)$, and any points \tilde{x} and \tilde{y} covering x and y respectively, there is a lifting \tilde{f} of f such that $\tilde{f}(\tilde{x}) = \tilde{y}$. So we have the following

Lemma 2.5. *Let $p : \widetilde{M} \rightarrow M$ be a characteristic covering between compact manifolds, and \tilde{w} be a lifting of an f -route w of a homeomorphism $f : M \rightarrow M$. Then there is a lifting \tilde{f} of f such that \tilde{w} is an \tilde{f} -route.*

3. SOME FACTS ON FUCHSIAN GROUPS

In this section, we give some facts on Fuchsian groups.

Let B be an orbifold. Recall that an orbifold in this paper means a compact 2-orbifold with singular points consisting of cone points, then we can assume $B = F(n_1, n_2, \dots, n_k)$, where the compact surface F denotes the underlying space of B and $n_i \geq 2$ denotes the cone point with cone angle $2\pi/n_i$, $i = 1, 2, \dots, k$. The fundamental group $\pi_1(B)$ is called a *Fuchsian group*. If H is a subgroup of the Fuchsian group $\pi_1(B)$, then H is also a Fuchsian group because there is a covering orbifold \tilde{B} of B such that $H \cong \pi_1(\tilde{B})$. Moreover, if \tilde{B} is also compact, then $\chi(\tilde{B}) = k\chi(B)$ and H has finite index k in $\pi_1(B)$, where $k > 0$ is the degree of the covering. In particular, if the orbifold B is hyperbolic, then B can be covered by a hyperbolic surface S , and $\pi_1(B)$ is an infinite Fuchsian group since $\pi_1(B)$ has a subgroup isomorphic to the infinite group $\pi_1(S)$. For more information on Fuchsian groups, see [JS, Chapter 2].

Lemma 3.1. (1) *Any infinite Fuchsian group with nontrivial center is either free abelian of rank ≤ 2 or isomorphic to the fundamental group of a Klein bottle.*

(2) *Any finite Fuchsian group is either cyclic or isomorphic to*

$$\langle a, b | a^{n_1} = b^{n_2} = (ab)^{n_3} = 1 \rangle, \quad \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} > 1, n_1, n_2, n_3 \geq 2,$$

which is the fundamental group of the closed orbifold $O = S^2(n_1, n_2, n_3)$ with $\chi(O) = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3} - 1 > 0$.

Proof. Conclusion (1) is from [JS, Proposition II.3.11], and Conclusion (2) can be verified by [He, Theorem 12.2] clearly. \square

For a group G and an element $g \in G$, let $C_G(g) = \{x \in G | xg = gx\}$ be the centralize of g in G , and $C(G) = \{x \in G | xg = gx, \forall g \in G\}$ the center of G . We have

Lemma 3.2. *Let B be a hyperbolic orbifold and $a \in \pi_1(B)$ be an element of infinite order. Then for any $i \neq 0$,*

$$C_{\pi_1(B)}(a^i) = C_{\pi_1(B)}(a) \cong \mathbb{Z}.$$

Proof. For any $i \neq 0$, $C_{\pi_1(B)}(a^i) \leq \pi_1(B)$ is an infinite Fuchsian group with nontrivial center since the infinite cyclic group $\langle a \rangle \leq C_{\pi_1(B)}(a^i)$. Then $C_{\pi_1(B)}(a^i)$ is either free abelian of rank ≤ 2 or the fundamental group of a Klein bottle by Lemma 3.1(1). Note that B is hyperbolic and $C_{\pi_1(B)}(a^i)$ is a subgroup of $\pi_1(B)$, then $C_{\pi_1(B)}(a^i)$ is neither the fundamental group of a Klein bottle nor the fundamental group of a torus because neither of them is hyperbolic. Therefore, $C_{\pi_1(B)}(a^i)$ is free cyclic, set $C_{\pi_1(B)}(a^i) = \langle c \rangle \cong \mathbb{Z}$. Since $a \in C_{\pi_1(B)}(a^i) = \langle c \rangle$, a is a power of c . Thus $\langle c \rangle \leq C_{\pi_1(B)}(a)$, i.e., $C_{\pi_1(B)}(a^i) \leq C_{\pi_1(B)}(a)$. Clearly, $C_{\pi_1(B)}(a) \leq C_{\pi_1(B)}(a^i)$. Thus $C_{\pi_1(B)}(a^i) = C_{\pi_1(B)}(a) \cong \mathbb{Z}$ for any $i \neq 0$. \square

Now we give two lemmas which are used in the following.

In group theory, a group G is called *metacyclic* if it contains a cyclic, normal subgroup N such that the quotient group G/N is also cyclic. Clearly, the rank of a metacyclic group is no more than 2. In particular, cyclic groups are metacyclic. For metacyclic groups, we have

Lemma 3.3. (1) *Any subgroup of a metacyclic group is also metacyclic.*

(2) *Let G be a group. If G has a cyclic normal subgroup T such that the quotient group G/T is metacyclic, then any subgroup of G has rank ≤ 3 .*

Proof. (1) Let G be a metacyclic group, i.e., there is a cyclic normal subgroup N such that G/N is also cyclic. Let H be any subgroup of G . Then $H \cap N$ is a cyclic normal subgroup of H , and $H/H \cap N \cong HN/N$ is also cyclic since $HN/N \leq G/N$. Thus H is metacyclic.

(2) Let H be any subgroup of G . Note that $H \cap T$ is a cyclic normal subgroup of H because T is cyclic normal in G . Then the quotient group $H/H \cap T \cong HT/T \leq G/T$ is metacyclic and $\text{rank}(H/H \cap T) \leq 2$ according to conclusion (1). Thus $\text{rank} H \leq 3$. \square

Lemma 3.4. *Suppose G is a group and $c \in G$ is an element of infinite order. If all the infinite-order elements are contained in the infinite cyclic group $\langle c \rangle$, then G is a metacyclic group. More precisely, either $G = \langle c \rangle \cong \mathbb{Z}$ or $G = \langle a, c | a^2 = 1, aca^{-1} = c^{-1} \rangle \cong \mathbb{Z}_2 * \mathbb{Z}_2$.*

Proof. If there are no nontrivial finite-order elements in G , then $G = \langle c \rangle \cong \mathbb{Z}$ clearly.

Now we assume that there is an element $a \in G$ of finite order $n \geq 2$. Consider the subgroup $H = \langle a, c \rangle \leq G$. Clearly aca^{-1} is also an element of infinite order, thus $aca^{-1} \in \langle c \rangle$ and $\langle c \rangle$ is a normal subgroup of H . Hence $aca^{-1} = c^i$ for some $i \neq 0$, and H is a metacyclic group. Then

$$c = a^n ca^{-n} = a^{n-1}(aca^{-1})a^{1-n} = a^{n-1}c^i a^{-n+1} = c^{i^n}.$$

Note that c is of infinite order, we have $i^n = 1$. Thus $i = 1$, if n is odd; $i = \pm 1$, if n is even. Namely, H is either

- (1) $\langle a, c | a^n = 1, aca^{-1} = c \rangle$, or
- (2) $\langle a, c | a^n = 1, aca^{-1} = c^{-1} \rangle$, n even.

However, case (1) is impossible. In fact, note that $ac \in H$ is of infinite order, thus $ac \in \langle c \rangle$. Then $a \in \langle c \rangle$ which contradicts to the assumption that $a \in G$ has finite order $n \geq 2$.

In case (2), note that $a^2 ca^{-2} = c$, we have an abelian subgroup $\langle a^2, c | a^n = 1, a^2 ca^{-2} = c \rangle$. It implies $a^2 c$ has infinite order. Thus $a^2 c \in \langle c \rangle$ and $a^2 \in \langle c \rangle$. Recall that a has finite order $n \geq 2$ and c has infinite order, then $n = 2$. It implies all the nontrivial finite-order elements in G must have order 2, and

$$H = \langle a, c | a^2 = 1, aca^{-1} = c^{-1} \rangle \cong \mathbb{Z}_2 * \mathbb{Z}_2.$$

To prove $G = H$, it suffices to prove $G \leq H$. Suppose there is another element $b \in G$ of order 2, then there is a subgroup $H' = \langle b, c | b^2 = 1, bcb^{-1} = c^{-1} \rangle$ according to the argument above. Thus $abcb^{-1}a^{-1} = c$, i.e., ab commutes with c . If abc has finite order $k > 0$, then $(ab)^k c^k = (abc)^k = 1$, i.e., $(ab)^k = c^{-k}$ has infinite order. Thus ab has infinite order and $ab \in \langle c \rangle$, it implies $b \in \langle a, c \rangle = H$. If abc has infinite order, then $abc \in \langle c \rangle$ which also implies $b \in \langle a, c \rangle = H$. Therefore, $G \leq H$ and the proof is finished. \square

Proposition 3.5. *Let H be a subgroup of $\pi_1(B)$ where B is a hyperbolic orbifold. If*

$$H^d := \{h^d | h \in H\} \subseteq \langle a \rangle \cong \mathbb{Z}$$

*where $d \in \mathbb{Z}_+$ is a positive integer and $a \in \pi_1(B)$ of infinite order. Then H is a metacyclic group, more precisely, H is either cyclic or isomorphic to $\mathbb{Z}_2 * \mathbb{Z}_2$.*

Proof. Note that B is a hyperbolic orbifold, namely, it can be covered by a hyperbolic surface S . Then $\pi_1(B)$ is an infinite Fuchsian group since it has a subgroup isomorphic to the infinite group $\pi_1(S)$.

If $H^d = \{1\}$, then H consists of elements of finite order. Thus H is a finite Fuchsian group by the fact that every infinite Fuchsian group has an element of infinite order (see [JS, Lemma II.3.9]). Therefore, H is either finite cyclic or isomorphic to the fundamental group $\pi_1(O)$ of a closed orbifold

O with $\chi(O) > 0$ by Conclusion (2) of Lemma 3.1. But the latter is impossible because B with $\chi(B) < 0$ can not be covered by O . Thus H is finite cyclic.

If $H^d \neq \{1\}$, then H contains some elements of infinite order. For any $h \in H$ of infinite order, there is an element $a^{i_h} \neq 1$ such that $h^d = a^{i_h}$ since $H^d \subseteq \langle a \rangle$. Then $h \in C_{\pi_1(B)}(a^{i_h})$. By Lemma 3.2, $C_{\pi_1(B)}(a^{i_h}) = C_{\pi_1(B)}(a) = \langle c \rangle$ for an infinite-order element c , thus $h \in \langle c \rangle$. Then all the infinite-order elements of H are contained in the infinite cyclic group $H \cap \langle c \rangle$. Therefore, H is either infinite cyclic or isomorphic to $\mathbb{Z}_2 * \mathbb{Z}_2$ by Lemma 3.4, and the proof is finished. \square

4. HOMEOMORPHISMS OF CIRCLE BUNDLES OVER SURFACES

In this section, let us consider the special case that M is an orientable circle bundle over an orientable hyperbolic surface S with fibration $p : M \rightarrow S$.

Lemma 4.1. *Suppose M is a compact orientable circle bundle over a compact orientable hyperbolic surface S . Then all fibers can be coherently oriented so that they present the same element of infinite order in the center of $\pi_1(M)$.*

Proof. This is clear from the presentation of $\pi_1(M)$ given in [He, Chapter 12]. \square

Let t be the element of $\pi_1(M)$ represented by the fiber. We call the infinite cyclic subgroup $\langle t \rangle$ the *fiber* of $\pi_1(M)$ associated to the fibration p , and f *preserves* (resp. *reverses*) the *fiber orientation* if $f_\pi(t) = t$ (resp. $f_\pi(t) = -t$).

Let S be a compact connected hyperbolic surface. A standard form of homeomorphisms on S is developed in [JG] with fine-tuned local behavior from the Thurston canonical map from [T].

Theorem T. *Let S be a compact connected hyperbolic surface. Every homeomorphism $f : S \rightarrow S$ is isotopic to a homeomorphism φ such that either*

- (1) φ is a periodic map, i.e., $\varphi^m = \text{id}$ for some $m \geq 1$, or equivalently, φ is an isometry with respect to some hyperbolic metric on S ; or
- (2) φ is a pseudo-Anosov map, i.e., there is a number $\lambda > 1$ and a pair of transverse measured foliations (\mathfrak{F}^s, μ^s) and (\mathfrak{F}^u, μ^u) such that $\varphi(\mathfrak{F}^s, \mu^s) = (\mathfrak{F}^s, \frac{1}{\lambda}\mu^s)$ and $\varphi(\mathfrak{F}^u, \mu^u) = (\mathfrak{F}^u, \lambda\mu^u)$; or
- (3) φ is a reducible map, i.e. there is a system of disjoint simple closed curves $\Gamma = \{\Gamma_1, \dots, \Gamma_k\}$ in $\text{int}S$ with the property below.
 - (a) Γ is invariant by φ (but the Γ_i 's may be permuted) and each component of $S \setminus \Gamma$ has negative Euler characteristic.
 - (b) Γ has a φ -invariant tubular neighborhood $\mathcal{N}(\Gamma)$ such that on each (not necessarily connected) φ -component of $S \setminus \mathcal{N}(\Gamma)$, φ satisfies (1) or (2).
 - (c) Γ is minimal among all systems satisfying (a) and (b).
 - (d) φ is in the standard form as defined in [JG, page 79].

The φ above will be called the *standard form* isotopic to S .

Lemma 4.2. *Let $f : S \rightarrow S$ be a homeomorphism of an orientable compact hyperbolic surface S and \mathbf{F}_w be a fixed point class corresponding to an f -route w . Then*

- (1) *If \mathbf{F}_w is inessential, then $\text{rankFix}(f_w) \leq 1$;*
- (2) *If f is orientation-reversing, then $\text{rankFix}(f_w) \leq 1$.*

Proof. Case (1) is clearly following from [JWZ, Theorem 1.1].

Now we consider case (2). Since the index and rank of fixed point classes are both homotopy invariants, via an isotopy, we may assume that f is in standard form. If the fixed point class \mathbf{F}_w is empty, then case (2) holds according to case (1). If \mathbf{F}_w is nonempty, it is connected since each

Nielsen path of S can be deformed (rel. endpoints) into $\text{Fix}f$ by [JG, Lemmas 1.2, 2.2 and 3.4]. Thus $\text{rankFix}(f_w) = \text{rank}\pi_1(\mathbf{F}_w)$. Then case (2) can be proved by examining the list given in [JG, Lemma 3.6]. \square

The lemma below is useful in the proof of Proposition 4.4.

Lemma 4.3. *Suppose $p : M \rightarrow S$ is a compact orientable circle bundle over a compact orientable hyperbolic surface S . Suppose $f : M \rightarrow M$ is a fiber-preserving homeomorphism that reverse the fiber orientation, and $f' : S \rightarrow S$ is the induced homeomorphism. Let w be an f -route. Then the f -fixed point class \mathbf{F}_w is essential if and only if the f' -fixed point class $\mathbf{F}_{p \circ w}$ is essential.*

Proof. Since the index of fixed point classes is a homotopy invariant, via a fiber-preserving homotopy, we can assume that the induced homeomorphism f' is in standard form by [JW, Lemma 2.7]. Then $\text{ind}(f, \mathbf{F}_w)$ equals either $\text{ind}(f', \mathbf{F}_{p \circ w})$ or $2\text{ind}(f', \mathbf{F}_{p \circ w})$ by [JW, Lemma 2.9]. Thus the conclusion holds. \square

Proposition 4.4. *Suppose $p : M \rightarrow S$ is a compact orientable circle bundle over an orientable hyperbolic surface S , $f : M \rightarrow M$ is a fiber-preserving homeomorphism that reverses the orientation of M , and $f' : S \rightarrow S$ is the induced homeomorphism of f . Let w be an f -route corresponding to an inessential fixed point class \mathbf{F}_w . Then*

- (1) $\text{Fix}(f'_{p \circ w})$ is trivial or the free cyclic group \mathbb{Z} ;
- (2) $\text{Fix}(f_w)$ is trivial or a free abelian group of rank ≤ 2 .

Proof. (1) Note that all fibers can be coherently oriented since M is a connected orientable circle bundle over an orientable surface by Lemma 4.1. So f is in one of the two cases below.

Case (i). f is fiber orientation-reversing and $f' : S \rightarrow S$ is orientation-preserving.

Since the f -fixed point class \mathbf{F}_w is inessential, the f' -fixed point class $\mathbf{F}_{p \circ w}$ is also inessential by Lemma 4.3. Then $\text{rankFix}(f'_{p \circ w}) \leq 1$ by Lemma 4.2(1).

Case (ii). f is fiber orientation-preserving and $f' : S \rightarrow S$ is orientation-reversing. Then $\text{rankFix}(f'_{p \circ w}) \leq 1$ by Lemma 4.2(2).

Hence conclusion (1) holds in both case (i) and case (ii) by the fact that every nontrivial element of an orientable surface group has infinite order.

(2) Let $x = w(0)$. Since $p \circ f = f' \circ p : M \rightarrow S$, there is the following commutative diagram on fundamental groups:

$$\begin{array}{ccc} \pi_1(M, x) & \xrightarrow{f_w} & \pi_1(M, x) \\ p_* \downarrow & & \downarrow p_* \\ \pi_1(S, p(x)) & \xrightarrow{f'_{p \circ w}} & \pi_1(S, p(x)) \end{array}$$

where

$$p_* : \pi_1(M, x) \rightarrow \pi_1(S, p(x)) \cong \pi_1(M, x) / \langle t \rangle$$

is the quotient map and $\langle t \rangle$ is the fiber of $\pi_1(M, x)$ associated with the fibration p . Hence

$$p_* \text{Fix}(f_w) \leq \text{Fix}(f'_{p \circ w}).$$

Since $\langle t \rangle$ is in the center of $\pi_1(M, x)$, and $\text{Fix}(f'_{p \circ w})$ is trivial or the infinite cyclic group \mathbb{Z} by conclusion (1), we have

$$\text{Fix}(f_w) \leq p_*^{-1} \text{Fix}(f'_{p \circ w}) \cong \text{Fix}(f'_{p \circ w}) \times \langle t \rangle,$$

which is isomorphic to the free abelian group \mathbb{Z} or $\mathbb{Z} \oplus \mathbb{Z}$. Hence conclusion (2) holds by the fact that every nontrivial subgroup of the free abelian group $\mathbb{Z} \oplus \mathbb{Z}$ is a free abelian group of rank ≤ 2 . \square

5. PROOF OF THEOREM 1.4

In this section, we give the proof of Theorem 1.4.

Let M be a compact orientable Seifert manifold with hyperbolic orbifold B_M , and $p : M \rightarrow B_M$ be the Seifert fibration. If τ is a regular fiber in this fibration, then

$$\langle t \rangle = \text{Im}(\pi_1(\tau, x) \rightarrow \pi_1(M, x))$$

is a infinite cyclic subgroup. We call $\langle t \rangle$ the *fiber* of $\pi_1(M, x)$ (associated with the fibration p). In fact, for each Seifert fibration of M , there exists a unique fiber of $\pi_1(M, x)$. It is well known that the cyclic subgroup $\langle t \rangle$ is normal in $\pi_1(M, x)$ and is center when M and B_M are both orientable. The Seifert fibration p induced a quotient homomorphism $p_* : \pi_1(M, x) \rightarrow \pi_1(M, x)/\langle t \rangle \cong \pi_1(B_M)$ on fundamental groups (see [JS, Chapter II, §4] or [He, Chapter 12]).

Proposition 5.1. *Let M be a compact orientable Seifert manifold with hyperbolic orbifold B_M , and $f : M \rightarrow M$ be an orientation-reversing homeomorphism. Suppose $f_w : \pi_1(M, x) \rightarrow \pi_1(M, x)$ is the automorphism induced by f , where w is an f -route with $x = w(0)$. If the fixed point class \mathbf{F}_w corresponding to w is inessential, then $\text{rankFix}(f_w) \leq 3$.*

To prove Proposition 5.1, we need the following useful lemma which is from [JWW, Theorem 3.11].

Lemma 5.2. *Suppose M is a compact orientable Seifert manifold and $p : M \rightarrow B_M$ is a Seifert fibration with hyperbolic orbifold B_M . Then any homeomorphism on M is isotopic to a fiber-preserving homeomorphism with respect to this fibration.*

Proof of Proposition 5.1. By Lemma 5.2, f can be isotopic to a fiber-preserving homeomorphism. Since the index $\text{ind}(f, \mathbf{F}_w)$ and $\text{rankFix}(f_w)$ of fixed point class are homotopy invariants (see Fact (homotopy invariance) in Section 2), we can assume that f is fiber-preserving and the base point x is in a regular fiber τ in the following.

Note that B_M is hyperbolic, then there is a finite covering $q : S \rightarrow B_M$ of orbifold such that S is a compact orientable hyperbolic surface by [S, Theorem 2.5]. The pull-back of the Seifert fibration $p : M \rightarrow B_M$ via q gives a finite covering manifold $q' : \widetilde{M} \rightarrow M$ with fibration $p' : \widetilde{M} \rightarrow B_{\widetilde{M}} = S$. After passing to further finite covering if necessary, we may assume that q' is a characteristic covering with finite sheets d . Since S is an orientable surface, the fibration p' is an orientable circle bundle over the orientable surface S . So there is the following commutative diagram:

$$\begin{array}{ccc} \widetilde{M} & \xrightarrow{q'} & M \\ p' \downarrow & & \downarrow p \\ S & \xrightarrow{q} & B_M \end{array}$$

Pick a point $\tilde{x} \in q'^{-1}(x)$ and a lifting \tilde{w} of w with $\tilde{w}(0) = \tilde{x}$. Since q' is a finite characteristic covering, following from Lemma 2.5, there is a lifting \tilde{f} of f such that \tilde{w} is an \tilde{f} -route. Let $\tilde{f}' : S \rightarrow S$ denote the induced homeomorphism of \tilde{f} on the orbifold S . Therefore, there is the following

commutative diagram on fundamental groups:

$$\begin{array}{ccccc}
 & & \pi_1(S, p'(x)) & \xrightarrow{\tilde{f}'_{p' \circ \tilde{w}}} & \pi_1(S, p'(x)) \\
 & \nearrow p'_* & \downarrow & & \nearrow p'_* \\
 \pi_1(\widetilde{M}, \tilde{x}) & \xrightarrow{\tilde{f}_{\tilde{w}}} & \pi_1(\widetilde{M}, \tilde{x}) & & \pi_1(\widetilde{M}, \tilde{x}) \\
 \downarrow q'_* & & \downarrow q'_* & & \downarrow q_* \\
 & \nearrow p_* & \pi_1(B_M) & \xrightarrow{\quad} & \pi_1(B_M) \\
 \pi_1(M, x) & \xrightarrow{f_w} & \pi_1(M, x) & & \pi_1(M, x) \\
 & \nwarrow p_* & \downarrow & & \nwarrow p_*
 \end{array}$$

Note that $f : M \rightarrow M$ is a fiber-preserving homeomorphism that reverse the orientation of M , then the lifting $\tilde{f} : \widetilde{M} \rightarrow \widetilde{M}$ is a fiber-preserving homeomorphism that reverse the orientation of \widetilde{M} . Moreover, since the f -fixed point class \mathbf{F}_w is inessential, the \tilde{f} -fixed point class $\mathbf{F}_{\tilde{w}}$ is also inessential by Lemma 2.4. Therefore, $\text{Fix}(\tilde{f}'_{p' \circ \tilde{w}})$ is trivial or the free cyclic group \mathbb{Z} according to conclusion (1) of Proposition 4.4. We claim that

Claim 5.3. *Let $H = p_*\text{Fix}(f_w) \subseteq \pi_1(B_M)$ and $H^d = \{h^d | h \in H\}$. Then*

$$H^d \subseteq q_*\text{Fix}(\tilde{f}'_{p' \circ \tilde{w}}) \leq \pi_1(B_M)$$

where $q_*\text{Fix}(\tilde{f}'_{p' \circ \tilde{w}})$ is trivial or the free cyclic group \mathbb{Z} .

Proof. Let $\alpha \in \text{Fix}(f_w) \leq \pi_1(M, x)$. Recall that $q' : \widetilde{M} \rightarrow M$ is a characteristic covering of sheets d , then

$$q'_* : \pi_1(\widetilde{M}, \tilde{x}) \rightarrow \pi_1(M, x)$$

is an injective homomorphism and the image $q'_*\pi_1(\widetilde{M}, \tilde{x})$ is a characteristic subgroup of $\pi_1(M, x)$ with index d . Hence the power

$$\alpha^d \in q'_*\pi_1(\widetilde{M}, \tilde{x}) \leq \pi_1(M, x),$$

furthermore, following from the commutative diagram, we have

$$\tilde{f}_{\tilde{w}}q'^{-1}(\alpha^d) = q'^{-1}f_w(\alpha^d) = q'^{-1}(\alpha^d),$$

and

$$\tilde{f}'_{p' \circ \tilde{w}}p'_*(q'^{-1}(\alpha^d)) = p'_*\tilde{f}_{\tilde{w}}(q'^{-1}(\alpha^d)) = p'_*q'^{-1}(\alpha^d).$$

Namely,

$$p'_*q'^{-1}(\alpha^d) \in \text{Fix}(\tilde{f}'_{p' \circ \tilde{w}}).$$

Then

$$(p_*(\alpha))^d = p_*(\alpha^d) = q_*p'_*q'^{-1}(\alpha^d) \in q_*\text{Fix}(\tilde{f}'_{p' \circ \tilde{w}}) \leq \pi_1(B_M).$$

Hence

$$H^d \subseteq q_*\text{Fix}(\tilde{f}'_{p' \circ \tilde{w}}) \leq \pi_1(B_M).$$

Note that q_* is induced by the covering q , then q_* is an injective homomorphism, and $q_*\text{Fix}(\tilde{f}'_{p' \circ \tilde{w}})$ is isomorphic to $\text{Fix}(\tilde{f}'_{p' \circ \tilde{w}})$ which is trivial or the free cyclic group \mathbb{Z} . Hence the Claim holds.

Now we will finish the proof of Proposition 5.1.

By Proposition 3.5 and Claim 5.3, $H = p_*\text{Fix}(f_w)$ is a metacyclic group. Since

$$p_* : \pi_1(M, x) \rightarrow \pi_1(M, x)/\langle t \rangle = \pi_1(B_M)$$

is the quotient map, we have $\text{Fix}(f_w) \leq p_*^{-1}(H)$ which is an extension of the metacyclic group H by the infinite cyclic group $\langle t \rangle$, namely, $\text{Fix}(f_w)$ is a subgroup of the group $p_*^{-1}(H)$ which has a cyclic normal subgroup $\langle t \rangle$ such that the quotient group H is metacyclic. So by Lemma 3.3, $\text{rankFix}(f_w) \leq 3$. \square

Now we give the proof of Theorem 1.4. Firstly, we have the following lemma on the ranks of Seifert manifold groups.

Lemma 5.4. *Suppose M is a compact orientable Seifert manifold with hyperbolic orbifold B_M . Then $\text{rank}\pi_1(M) \geq 2$.*

Proof. It is clear from [Z1, Proposition 4.3] which gives a description on the ranks of Seifert manifold groups. \square

Proof of Theorem 1.4. Let $f_\pi = f_w : \pi_1(M, x) \rightarrow \pi_1(M, x)$ be induced by some f -route w with $x = w(0)$.

If \mathbf{F}_w is essential, then $\text{rankFix}(f_w) < 2\text{rank}\pi_1(M)$ by Proposition 1.7.

If \mathbf{F}_w is inessential, then $\text{rankFix}(f_w) \leq 3$ by Proposition 5.1. Recall that the orbifold B_M of M is hyperbolic, then $\text{rank}\pi_1(M) \geq 2$ according to Lemma 5.4. Therefore, we have

$$\text{rankFix}(f_w) < 2\text{rank}\pi_1(M).$$

The proof is finished. \square

6. EXAMPLES AND QUESTIONS

Now we give some examples and questions.

Example 6.1. Let S_n be a closed orientable surface of genus $n \geq 2$. Define an orientation-reversing homeomorphism f as follows:

$$f = f_1 \times f_2 : S_n \times S^1 \rightarrow S_n \times S^1,$$

where $f_1 : S_n \rightarrow S_n$ is a reflection on a simple closed curve γ , and $f_2 : S^1 \rightarrow S^1$ is a rotation. Then all the fixed point classes of f are inessential, and f induces an automorphism f_π of $\pi_1(S_n \times S^1)$ such that

$$\text{Fix}(f_\pi) = \pi_1(\gamma \times S^1) \cong \mathbb{Z} \oplus \mathbb{Z}.$$

Namely, there is an inessential fixed point class which has $\text{rankFix}(f_\pi) = 2$.

Question 6.2. Is there an orientation-reversing homeomorphism f of a Seifert manifold M whose inessential fixed point class has $\text{rankFix}(f_\pi) = 3$? Namely, is the bound 3 in Proposition 5.1 sharp?

REFERENCES

- [BH] M. Bestvina and M. Handel, *Train tracks and automorphisms of free groups*, Annals of Mathematics, 135 (1992), 1–51.
- [He] J. Hempel, *3-manifolds*, Annals of Math. Studies 86, Princeton University Press, 1976.
- [J] B. Jiang, *Lectures on Nielsen Fixed Point Theory*, Contemporary Mathematics vol. 14, American Mathematical Society, Providence (1983).
- [JG] B. Jiang and J. Guo, *Fixed points of surface diffeomorphisms*, Pacific Journal of Mathematics, 160 No.1 (1993), 67–89.
- [JS] W. Jaco and P. Shalen, *Seifert fibered spaces in 3-manifolds*, Memoirs of the American Mathematical Society; V.21, No.220, Providence (1979).
- [JW] B. Jiang and S. Wang, *Lefschetz numbers and Nielsen numbers for homeomorphisms on aspherical manifolds*, Topology Hawaii, 1990, World Sci. Publ., River Edge, NJ, 1992, 119–136.
- [JWW] B. Jiang, S. Wang and Y. Wu, *3-manifolds and the realization of Nielsen number*, Comm. Anal. Geom. Vol. 9(2001), NO.4, 825–877.
- [JWZ] B. Jiang, S. Wang and Q. Zhang, *Bounds for fixed points and fixed subgroups on surfaces and graphs*, Alg. Geom. Topology, 11(2011), 2297–2318.

- [LW] J. Lin and S. Wang, *Fixed subgroups of automorphisms of hyperbolic 3-manifold groups*, Topology Appl., 173(2014), 209–218 .
- [S] P. Scott, *The geometries of 3-manifolds*, Bull. London Math. Soc. 15 (1983), 401–487.
- [T] W.P. Thurston, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc, 19 (1988), 417–431.
- [WZ] J. Wu and Q. Zhang, *The group fixed by a family of endomorphisms of a surface group*, J. Algebra, 417(2014), 412–432 (2014)
- [Z1] Q. Zhang, *Bounds for fixed points on Seifert manifolds*, Topology and its Applications, 159(15) (2012), 3263–3273.
- [Z2] Q. Zhang, *Bounds for fixed points on hyperbolic 3-manifolds*, Topology Appl., 164(2014), 182–189.

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN 710049, CHINA
E-mail address: `zhangq.math@mail.xjtu.edu.cn`